arXiv:hep-th/0105112v2 19 Jun 2001

SOGANG-HEP 281/01
hep-th/0105112

Quasinormal Modes and Choptuik Scaling in the Near Extremal Reissner-Nordström Black Hole

Won Tae Kim* and John J. Oh†

Department of Physics and Basic Science Research Institute,
Sogang University, C.P.O. Box 1142, Seoul 100-611, Korea

(February 1, 2008)

Abstract

We study quasinormal modes of massless scalar and fermion fields in the near extremal Reissner-Nordström black hole, and relate them to Choptuik scaling form following a recently proposed analytic approach. For both massless cases, quasinormal modes are shown to be proportional to the black hole horizon and the Hawking temperature, and the critical exponents are the same, although for the fermionic case there are two possible discrete quasinormal modes. In addition, the critical exponent of the massive boson is also equivalent to that of the massless case. Finally, we discuss quasinormal modes and critical exponents in the other models, and obtain some different critical exponents between massless boson and massive one.

PACS : 04.62.+v, 04.70.-s

Typeset using REVTEX

---

*electronic address:wtkim@ccs.sogang.ac.kr

†electronic address:john5@string.sogang.ac.kr

One of the most exciting studies in gravitation is the gravitational collapse of infalling matters. When smooth initial data are slightly greater than those of critical ones, then the black hole formation appears, otherwise the matter fields eventually scattered below the critical values of initial data [1]. The dynamical behaviors are quite same and indistinguishable after certain time, and then the final state of the collapsing matter is determined by the initial data [2].

Since Bañados, Teitelboim, and Zanelli(BTZ) have found a nontrivial black hole solution in (2+1)-dimensional anti-de Sitter(AdS) spacetime which has a negative constant scalar curvature [3], it has been widely studied in diverse branches of theoretical physics, including gravity and string theory. Recently, in connection with a critical behavior of black hole formation in (2+1)-dimensional AdS spacetimes, several works have been done not only in numerical ways but also analytical ones. For a massless scalar field, critical behavior has been studied in Ref. [4] by supposing a scaling relation of an AdS black hole,

$$M = K(p - p^*)^{2\gamma}, \tag{1}$$

where the critical exponent yields $\gamma = 1.2 \pm 0.05$ by some numerical calculations. An analytical approach [5] gives an exact critical solution for a scalar field, which is compatible with the result obtained in Ref. [4]. By a different numerical way, the critical collapse of scalar field in (2+1)-AdS spacetimes has been studied and obtained the critical exponent for a black hole formation, $\gamma = 0.81$ [6]. In the case of dust-shell collapse, the exponent $\gamma = 1/2$ has been obtained [7]. Especially the BTZ black hole formation has been studied in terms of group generators of AdS$_3$ isometries using Gott time machine condition, and obtained a critical exponent $\gamma = 1/2$ for two colliding particles [8].

On the other hand, a few works on quasinormal modes of black holes show that quasinormal frequencies may be related to the critical exponents in some special spacetime geometries. The quasinormal modes of minimally coupled scalar fields in AdS-Schwarzschild black hole have been studied in three [9], four [10,11], five, and seven dimensions [10]. And a conjecture between quasinormal frequencies and a critical exponent, $\omega \sim r_H/\gamma$, has been given [10], and analytically studied in BTZ black hole background for a massive scalar field in Ref. [12].

In this paper, we shall study some field equations on the near extremal RN black hole background which is effectively described by AdS_2 black holes by assuming a spherical symmetric dimensional reduction. Quasinormal modes of massless scalar and fermionic fields in the near extremal RN black hole will be calculated following Ref. [12]. For both cases, quasinormal modes are shown to be proportional to the black hole horizon and the Hawking temperature, and the critical exponents γ_b, γ_f for boson and fermionic respectively are all the same, $\gamma_b = \gamma_f = 1/2$. Furthermore, the critical exponent of the massive boson is calculated, which turns out to be equivalent to that of the massless case. Finally, we shall discuss quasinormal modes and critical exponents in the Jackiw-Teitelboim(JT) model [13], which gives some different critical exponents between massless boson and massive one.

We start with the Einstein-Maxwell system with a classical matter in four dimensions, which is given by

$$S_{\text{total}} = S_{\text{EM}} + S_{\text{M}} = \frac{1}{16\pi G}\int d^4x\sqrt{-g_{(4)}}\left[R_{(4)} - \frac{1}{4}F^{\mu\nu}F_{\mu\nu}\right] + S_{\text{M}}, \tag{2}$$

where F is a field strength satisfying $F = dA$ and S_M is a matter action. The Einstein-Maxwell action S_{EM} can be reduced to the two-dimensional action,

$$S = \frac{1}{2\pi}\int dx^2\sqrt{-g}e^{-2\phi}\left[R + 2(\nabla\phi)^2 + 2\lambda^2 e^{2\phi} - \frac{1}{4}F^2\right], \tag{3}$$

by assuming the spherically symmetric metric ansatz, $(ds)^2_{(4)} = (ds)^2 + (e^{-2\phi}/\lambda^2)d\Omega^2_{(2)}$ and $4\lambda^2 G = 2\pi$. The equation of motion for the action is exactly soluble and its solution is described by the two-dimensional charged Schwarzschild black hole,

$$(ds)^2 = -\left(1 - \frac{2GM}{x} + \frac{Q^2G^2}{x^2}\right)d^2t + \left(1 - \frac{2GM}{x} + \frac{Q^2G^2}{x^2}\right)^{-1}d^2x, \tag{4}$$

with the dilaton and gauge field solutions,

$$\begin{aligned} \frac{e^{-2\phi}}{\lambda^2} &= x^2, \\ e^{-2\phi}F^{\mu\nu} &= \pi Q\epsilon^{\mu\nu}, \end{aligned} \tag{5}$$

where Q is a electric charge of the black hole. In terms of the inner and outer horizons $x_\pm = GM \pm G\sqrt{M^2 - Q^2}$, Eq. (4) can be rewritten as

$$(ds)^2 = -\frac{(x - x_+)(x - x_-)}{x^2} d^2t + \frac{x^2 d^2 x}{(x - x_+)(x - x_-)}. \tag{6}$$

Then the Hawking temperature is

$$T_H = \frac{\kappa_+}{4\pi} = \frac{(x_+ - x_-)}{4\pi x_+^2}, \tag{7}$$

where κ_+ is a surface gravity at the event horizon.

We are now interested in the near extremal case of the metric (6). After performing the coordinate change such as $x = G\tilde{x} - 2\pi G^2 Q^2 T_H + x_+$ and $t = 1/G\tilde{t}$, and we take a near extremal limit $x_+ \to x_-$ under the fixed T_H and $G \to 0$ [14], then the metric (6) can be written as an AdS$_2$ black hole form,

$$(ds)^2 = -\left(-\mu^2 + \frac{\tilde{x}^2}{x_+^2}\right) d^2\tilde{t} + \left(-\mu^2 + \frac{\tilde{x}^2}{x_+^2}\right)^{-1} d^2\tilde{x}, \tag{8}$$

where $\mu^2 = (2\pi Q T_H)^2$. By performing one more coordinate transformation such as $\tilde{x} = (2x_+ G r - x_+^2)/(2x_+ G)$, and taking limits as $G \to 0$ and $x_+ \to x_-$ under the fixed Hawking temperature, we obtain the JT model [13] upto the leading order of a dilaton field,

$$\begin{aligned} (ds)^2 &= -\left(-\mu^2 + \frac{r^2}{x_+^2}\right) d^2t + \left(-\mu^2 + \frac{r^2}{x_+^2}\right)^{-1} d^2r + \mathcal{O}(r^3), \\ e^{-2\phi} &= \pi x_+ r. \end{aligned} \tag{9}$$

So, the near extremal RN black hole can be effectively described by the JT model [15]. Note that we assume $G \to 0$, and this gives a large mass black hole for a finite outer horizon x_+, since $GM \sim x_+ \sim GQ$ in the near extremal limit. Therefore, as the black hole mass becomes large, the black hole charge also becomes large. In Eq. (9), since $\mu \sim 1/G^2 x_+$ as $G \to 0$, the horizon of AdS black hole becomes large in the near extremal limit. The scalar curvature for Eq. (4) is

$$R = \frac{6Q^2G^2 - 4GMx}{x^4}, \tag{10}$$

and in an appropriate limit of the near extremal black hole, the given scalar curvature approaches to the constant, $R \propto -1/x_+^2$.

Next, we shall study quasinormal modes of some matter fields in this near extremal RN black hole in the spherical symmetric reduction, and relate quasinormal modes to the critical exponent of scaling law for the black hole formation. So the massless bosonic and fermionic fields in four dimensions are first considered, which is effectively described by the following two-dimensional action,

$$S_M = \frac{1}{2\pi} \int dx^2 \sqrt{-g} e^{-2\xi\phi} \left[-\frac{1}{2} (\nabla f)^2 \right], \tag{11}$$

by taking only s-wave sector of matter fields. Note that for $\xi = 1$, it describes a bosonic field in four dimensions, while the fermionic case is for $\xi = 0$ which is written after bosonization [16]. The equation of motion from the action (11),

$$\Box f - 2\xi \partial_\mu \phi \partial^\mu f = 0, \tag{12}$$

can be solved under the dilatonic AdS$_2$ black hole background (9) This metric has a new horizon at $r = r_H \equiv x_+ \mu$, and a new Hawking temperature $T_H^{\rm AdS}$ is related to the original temperature as $T_H^{\rm AdS} = (Q/x_+) T_H$.

In fact, quasinormal modes in AdS geometry has been defined in Refs. [10,12]. In the near horizon limit, only ingoing waves can be allowed while some fields should vanish at the AdS boundary since the potential diverges at infinity. This is a prominent feature of quasinormal modes in AdS geometry which differs from those in asymptotically flat one. So there exists a discrete set of quasinormal modes ω with a negative imaginary part [10]. In the two-dimensional system with an AdS geometry, we expect the solution of wave equation can be exactly solved and will be a form of a hypergeometric function [17].

First, we consider the bosonic case of $\xi = 1$ to solve the wave equation (12). Using the background metric (9) and the separation of variables $f(r,t) = R(r) e^{-i\omega t}$, the equation of motion can be written as

$$\left(r^2 - r_H^2 \right) \partial_r^2 R(r) + \frac{1}{r} \left(3r^2 - r_H^2 \right) \partial_r R(r) + \frac{\omega^2 x_+^4}{(r^2 - r_H^2)} R(r) = 0, \tag{13}$$

where $r_H = \mu x_+$. By means of the coordinate change $z = (r^2 - r_H^2)/r^2$, Eq. (13) becomes

$$z(1-z)\partial_z^2 R(z) + (1-z)\partial_z R(z) + \frac{A}{z} R(z) = 0, \tag{14}$$

where $A = \omega^2 x_+^4 / 4 r_H^2$. Note that a new variable z spans from 0 to 1 as r goes from r_H to infinity. To remove a singular point at $z = 0$, we set $R(r) = z^\alpha g(z)$. Then the equation of motion can be solved exactly, which yields a solution of hypergeometric functions $F(a, b, c; z)$ and $z^{1-c} F(a - c + 1, b - c + 1, 2 - c; z)$, and α^2 is determined as $-A$, where $a = b = \alpha$, and $c = 1 + 2\alpha$. So, we have a general solution from Eq. (13),

$$R(z) = C_{\rm out} z^\alpha F(\alpha, \alpha, 1 + 2\alpha; z) + C_{\rm in} z^{-\alpha} F(-\alpha, -\alpha, 1 - 2\alpha; z), \tag{15}$$

where $C_{\rm in}$ and $C_{\rm out}$ are ingoing and outgoing coefficients, respectively, and $\alpha = i\omega x_+^2 / 2 r_H$. We take a plus sign of α due to $\alpha \to -\alpha$ symmetry for convenience.

Applying the quasinormal mode condition, $C_{\rm out} = 0$ at the horizon as $z \to 0$, and we have only ingoing near horizon solution,

$$R(z) = C_{\rm in} z^{-\alpha} F(-\alpha, -\alpha, 1 - 2\alpha; z). \tag{16}$$

To give a boundary condition at the AdS boundary, we take a transformation of $z \to 1 - z$. In this case, since we have $c = a + b + m$ with $m = 1$, we have to use the following transformation rule [18],

$$\begin{aligned} F(a, b, a + b + m; z) = {} & \frac{\Gamma(m)\Gamma(a+b+m)}{\Gamma(a+m)\Gamma(b+m)} \sum_{n=0}^{m-1} \frac{(a)_n (b)_n}{n!(1-m)_n} (1-z)^n \\ & - \frac{\Gamma(a+b+m)}{\Gamma(a)\Gamma(b)} (z-1)^m \sum_{n=0}^{\infty} \frac{(a+m)_n (b+m)_n}{n!(n+m)!} (1-z)^n \\ & \times \left[\ln(1-z) - \psi(n+1) - \psi(n+m+1) + \psi(a+n+m) + \psi(b+n+m) \right]. \end{aligned} \tag{17}$$

To vanish the solution at infinity($z \to 1$), we take a condition of either $a + 1 = -n$ or $b + 1 = -n$. It gives a discrete quasinormal frequency ω as

$$\omega = -i \frac{2 r_H}{x_+^2} (n+1), \tag{18}$$

so that the critical exponent for the massless boson case is regarded as $\gamma_b^{(0)} = 1/2$. In terms of the Hawking temperature, Eq. (18) is written as

$$\omega = -i4\pi \frac{Q}{x_+} T_H(n+1) = -i4\pi T_H^{\rm AdS}(n+1). \tag{19}$$

So this quasinormal mode is proportional to the horizon r_H, and the Hawking temperature $T_H^{\rm AdS}$ of the AdS black hole [10].

In Eq. (18), the discretized quasinormal frequency has a negative imaginary part. The most general proof of this argument is given in Ref. [10]. It can be realized by assuming the field equation under the gravitational background with the boundary condition at the asymptotic region. By using the metric in terms of Eddington-Finkelstein coordinates, $v = t + r_* = t + r/g(r)$ as

$$(ds)^2 = -g(r)d^2v + 2dvdr, \tag{20}$$

where $g(r) = (\mu^2 - r^2/x_+^2)$ and performing separation of variables, $f(v,r) = R(r)e^{-i\omega v}$ with conditions $g(r_H) = 0$ and $R(\infty) = 0$, the useful relation is given as [10]

$$\int_{r_H}^{\infty} dr \left[g(r)|\frac{dR}{dr}|^2 \right] = -\frac{|\omega|^2 |R(r_H)|^2}{{\rm Im}\omega}. \tag{21}$$

Therefore, quasinormal frequency has a negative imaginary part since $g(r)$ is positive definite outside the horizon.

Now we consider the fermionic case of $\xi = 0$. In this case, we have somewhat different type of scalar field equation from Eq. (13) since there does not exist a nontrivial dilaton coupling in the field equation, which is given as

$$(r^2 - r_H^2)\partial_r^2 R(r) + 2r\partial_r R(r) + \frac{\omega^2 x_+^4}{(r^2 - r_H^2)} R(r) = 0. \tag{22}$$

Using the coordinate transformation $z = (r^2 - r_H^2)/r^2$ and setting $R(z) = z^\alpha g(z)$ to remove a singularity at $z = 0$, the wave equation becomes

$$z(1-z)\partial_z^2 g(z) + \frac{1}{2}\left[2 + 4\alpha - (3 + 4\alpha)z\right]\partial_z g(z) - \frac{1}{2}\alpha(2\alpha + 1)g(z) = 0, \tag{23}$$

where α^2 is determined as $-A = -\omega^2 x_+^4/4r_H^2$. The solution of Eq. (23) is given as

$$R(z) = D_{\rm in} z^{-\alpha} F\left(\frac{1}{2} - \alpha, -\alpha, 1 - 2\alpha; z\right) + D_{\rm out} z^{\alpha} F\left(\alpha, \frac{1}{2} + \alpha, 1 + 2\alpha; z\right), \tag{24}$$

where $D_{\rm in}$ and $D_{\rm out}$ are ingoing and outgoing coefficients, respectively, and $\alpha = i\omega x_+^2/2r_H$. From the definition of quasinormal modes, we set $D_{\rm out} = 0$. For the present case, one should use a different transformation rule of $z \to 1 - z$ [18],

$$\begin{aligned} F(a, b, c; z) = {} & \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F(a, b, a+b-c+1; 1-z) \\ & + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} F(c-a, c-b, c-a-b+1; 1-z), \end{aligned} \tag{25}$$

where $a = 1/2 - \alpha$, $b = -\alpha$, and $c = 1 - 2\alpha$. To vanish the solution as $z \to 1$ gives a condition either $c - a = -n$ or $c - b = -n$. This yields two types of discrete quasinormal frequencies as

$$\omega = -i\frac{2r_H}{x_+^2}\left(n + \frac{1}{2}\right), \tag{26}$$

or

$$\omega = -i\frac{2r_H}{x_+^2}(n+1). \tag{27}$$

As for the critical exponent for the massless fermionic case, it yields $\gamma_f^{(0)} = 1/2$, which is the same value with that of the bosonic case.

Now let us consider that a massive scalar field in the s-wave sector. It can be reduced to the two-dimensional massive scalar field equation in the dilatonic AdS black hole background,

$$\Box f - 2\partial_\mu \phi \partial^\mu f - m^2 f = 0. \tag{28}$$

Using the separation of variables $f(r,t) = R(r)e^{-i\omega t}$ and coordinate change $z = (r^2 - r_H^2)/r^2$ for $0 \le z \le 1$, the wave equation becomes

$$z(1-z)\partial_z^2 R(z) + (1-z)\partial_z R(z) + \left[\frac{\omega^2 x_+^4}{4r_H}\left(\frac{1}{z}\right) - \frac{m^2 x_+^2}{1-z}\right] R(z) = 0. \tag{29}$$

To remove two singular points at $z = 0$ and $z = 1$, one can redefine $R(z) = z^{\alpha}(1-z)^{\beta} g(z)$, Eq. (29) becomes a standard form of a differential equation satisfying a hypergeometric function,

$$z(1-z)\partial_z^2 g(z) + \left[1 + 2\alpha - (1 + 2\alpha + 2\beta) z\right] \partial_z g(z) - (\alpha+\beta)^2 g(z) = 0, \tag{30}$$

where $\alpha = i\omega x_+^2/2r_H$ and $\beta = (1 - \sqrt{1 + 4m^2 x_+^2})/2$. The general solution of Eq. (30) is

$$\begin{aligned} R(z) &= E_{\rm out} z^{\alpha}(1-z)^{\beta} F(\alpha+\beta, \alpha+\beta, 1+2\alpha; z) \\ &\quad + E_{\rm in} z^{-\alpha}(1-z)^{\beta} F(-\alpha+\beta, -\alpha+\beta, 1-2\alpha; z), \end{aligned} \tag{31}$$

and $E_{\rm out} = 0$ by the definition of quasinormal modes. At this stage, we use the transformation rule (25) to make the solution vanish at infinity, and this gives a quasinormal frequency

$$\omega = -i\frac{2r_H}{x_+^2}\left(n + \frac{1}{2} + \frac{1}{2}\sqrt{1 + 4m^2 x_+^2}\right), \tag{32}$$

and it gives the critical exponent for the massive boson, $\gamma_b^{(m)} = 1/2$.

We have studied quasinormal modes and critical exponents of classical matters including massless boson and fermion, and massive bosonic field of s-wave sector in the near extremal RN black hole. Quasinormal frequencies are obtained by solving the wave equation from the definition of quasinormal modes. Following the relation $\omega \sim r_H/\gamma$ in Ref. [12], as a result, the critical exponents could be obtained as $\gamma_b^{(0)} = \gamma_f^{(0)} = \gamma_b^{(m)} = 1/2$ for massless scalar, massless fermion, and massive scalar fields.

Note that the charge scaling relation was not derived here because the black hole charge has been regarded as a fixed value in order to use AdS geometry and the useful relation to quasinormal modes. On the other hand, the charge scaling and its universality in critical collapse of charged scalar fields have been studied in Ref. [19], and a mass scaling, $M \sim (p - p_*)^{\gamma}$, and a new scaling with respect to the black hole charge, $Q \sim (p - p_*)^{\delta}$, have been obtained where the corresponding exponents, $\gamma = 0.374 \pm 0.001$ and $\delta = 0.883 \pm 0.007$, respectively. However, our approach is valid only for large black holes, since the RN black hole can be described by the dilatonic AdS$_2$ black hole which behaves as a large black hole in the near extremal limit. For these large black holes, it can be shown that quasinormal modes scale with the black hole temperature $T_H^{\rm AdS}$. These will decay exponentially, which translates into a time scale for the approach to the thermal equilibrium. Time scale is given by the imaginary part of the lowest quasinormal mode $\tau = 1/\omega_I$, where $\omega = \omega_R - i\omega_I$. These time scales have universal values in that all scalar fields will decay at these rates [10]. For our cases, the time scale of the lowest quasinormal mode can be obtained for $n = 0$ and inversely proportional to the Hawking temperature,

$$\tau \sim \frac{x_+^2}{r_H} = \frac{1}{T_H^{\rm AdS}}. \tag{33}$$

This means that the more black hole radiates, the faster it decays and approaches to the thermal equilibrium. For a massless bosonic case($\xi = 1$), we have a time scale $\tau_b^{(0)} = 1/4\pi T_H^{\rm AdS}$ while the time scale of a massive boson is given as

$$\tau_b^{(m)} = \frac{1}{2\pi T_H^{\rm AdS}} \left(1 + \sqrt{1 + 4m^2 x_+^2} \right)^{-1}. \tag{34}$$

Note that the fluctuated black hole from perturbations of the massive boson will go to the thermal equilibrium faster than those of the massless bosonic case since $\tau_b^{(0)} > \tau_b^{(m)}$. For the fermionic case, two kinds of time scale, $\tau_f^{(0)} = 1/4\pi T_H^{\rm AdS}$ and $\tau_f^{(0)} = 1/2\pi T_H^{\rm AdS}$, are possible due to Eqs. (26) and (27).

On the other hand, one might be interested in the critical exponents of two-dimensional scalar fields instead of the above dimensionally reduced four-dimensional ones on this dilatonic AdS$_2$ black hole background of the JT model [13], which is just the two-dimensional version of the initial work done in the BTZ black hole [12]. For the case of the two-dimensional massless boson, it is just what we have already done as far as we remove the dilaton coupling in Eq. (12), which corresponds to the equation of motion for the spherical symmetric fermion in the four dimensions. Therefore, the relation for the massless field in two dimensions is $\omega = -i2r_H/x_+^2(n+1)$ and $\gamma_{\rm massless}^{\rm AdS_2} = 1/2$. Then, what about the massive scalar field in the AdS$_2$ black hole background? The above similar calculations can be straightforwardly done, by dropping the dilaton coupling in Eq. (12) and adding mass term. Performing some change of variables as $z = (r - r_H)/(r + r_H)$ and imposing the above quasinormal condition, the following relation is obtained as $\omega = -ir_H/x_+^2(n+1)$. It gives interestingly twice value compared to that of the massless case, $\gamma^{\rm AdS_2}_{\rm massive} = 1$. This is more or less surprising in that the critical exponent of the massive scalar field is different from that of the massless one in the $\rm AdS_2$ black hole, whereas this is not the case for the BTZ black hole i.e., the critical exponents are the same, $\gamma^{\rm BTZ}_{\rm massive} = \gamma^{\rm BTZ}_{\rm massless} = 1/2$ [12].

The final comment is that we have not justified the scaling relation (1). Of course, in the three-dimensional BTZ case, it is proved in the numerical way in Refs. [4]. So, it would be interesting to find out how to obtain the squared scaling relation in two-dimensional AdS.

Acknowledgments

This work was supported by grant No. 2000-2-11100-002-5 from the Basic Research Program of the Korea Science and Engineering Foundation. We would like to thank to P. P. Jung for helpful discussions.

REFERENCES

[1] M. W. Choptuik, *Universality and Scaling in Gravitational Collapse of a Massless Scalar Field*, Phys. Rev. Lett. **70** (1993) 9.

[2] For a recent review, C. Gundlach, *Critical phenomena in gravitational collapse*, `[gr-qc/0001046]`.

[3] M. Bañados, C. Teitelboim, and J. Zanelli, *The Black Hole in Three-dimensional Spacetime*, Phys. Rev. Lett. **69** (1992) 1849, `[hep-th/9204099]`.

[4] F. Pretorius and M. W. Choptuik, Phys. Rev. D62 (2000) 124012, `[gr-qc/0007008]`.

[5] D. Garfinkle, *An exact solution for 2+1 dimensional critical collapse*, Phys. Rev. **D63** (2001) 044007, `[gr-qc/0008023]`.

[6] V. Husain and M. Olivier, *Scalar field collapse in three-dimensional AdS spacetime*, Class. Quant. Grav. **18** (2001) L1, `[gr-qc/0008060]`.

[7] Y. Peleg and A. R. Steif, *Phase Transition for Gravitationally Collapsing Dust Shells in (2+1)-dimensions*, Phys. Rev. **D51** (1995) 3992, `[gr-qc/9412023]`.

[8] D. Birmingham and S. Sen, *Gott Time Machines, BTZ Black Hole Formation, and Choptuik Scaling*, Phys. Rev. Lett. **84** (2000) 1074, `[hep-th/9908150]`.

[9] V. Cardoso and J. P. S. Lemos, *Scalar, Electromagnetic and Weyl Perturbations of BTZ black holes: Quasinormal Modes*, `[gr-qc/0101052]`.

[10] G. Horowitz and V. E. Hubeny, *Quasinormal modes of AdS black holes and the approach to thermal equilibrium*, Phys. Rev. **D62** (2000) 024027, `[hep-th/9909056]`.

[11] B. Wang, C.-Y. Lin, and E. Abdalla, *Quasinormal modes of Reissner-Nordstrom Anti-de Sitter black holes*, Phys. Lett. **B481** (2000) 79 ; J.-M. Zhu, B. Wang, and E. Abdalla, *Object picture of quasinormal ringing on the background of small Schwarzschild anti-de Sitter black holes*, Phys. Rev. **D63** (2001) 124004.

[12] D. Birmingham, *Choptuik Scaling and Quasinormal Modes in the AdS/CFT Correspondence*, `[hep-th/0101194]`.

[13] R. Jackiw, in Quantum Theory of Gravity, edited by S. M. Christensen (Hilger, Bristol, 1984); C. Teitelboim, in Quantum Theory of Gravity, edited by S. M. Christensen (Hilger, Bristol, 1984).

[14] M. Spradlin and A. Strominger, *Vacuum States for AdS$_2$ Black Holes*, J. High Energy Phys. **9911** (1999) 021, `[hep-th/9904143]`.

[15] A. Fabbri, D. J. Navarro, and J. Navarro-Salas, *Quantum evolution of near-extremal Reissner-Nordström black holes*, Nucl. Phys. **B595** (2001) 381, `[hep-th/0006035]`.

[16] M. G. Alford and A. Strominger, *S Wave Scattering of Charged Fermions by a Magnetic Black Hole*, Phys. Rev. Lett. **69** (1992) 563, `[hep-th/9202075]`.

[17] W. T. Kim and J. J. Oh, *Dilaton Driven Hawking Radiation in AdS$_2$ Black Hole*, Phys. Lett. **B461** (1999) 189, `[hep-th/9905007]`.

[18] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, Dover Publication Inc., New York, ninth printing, (1970).

[19] C. Gundlach and J. M. Martin-Garcia, *Charge scaling and universality in critical collapse*, Phys. Rev. **D54** (1996) 7353, `[gr-qc/9606072]`.